

A MEDICAL SOLUTION, A METHOD FOR PRODUCING SAID MEDICAL  
SOLUTION AND USE THEREOF

Technical field of the present invention

The present invention concerns a medical solution, a method for producing said medical solution, a multi-  
5 compartment bag containing the medical solution and the use thereof.

Background of the invention

Medical solutions like dialysis solutions for  
10 hemodialysis, hemofiltration, hemodiafiltration, peritoneal dialysis, dialysis within renal intensive care and liquids for substitution or infusion normally contain a buffering substance. Often used buffers are acetate and lactate buffers and these buffers are within the human  
15 body metabolized into bicarbonate. Thus, the most physiological buffer in medical solutions would be bicarbonate.

However, the use of bicarbonate as a buffer is more complicated than the use of acetate and lactate for two  
20 reasons: First bicarbonate easily precipitates with one of the essential elements in dialysis fluids, viz. calcium, to form calcium carbonate, and second bicarbonate solutions emit carbon dioxide and are thus unstable.

25 One way to get around the precipitation problem is to separate bicarbonate and calcium in two different containers and then mix them just before use, but the problem with the emitted carbon dioxide still remains.

If carbon dioxide leaves the bicarbonate solution  
30 the result is an increase of pH up to 9-10,5 depending on the original bicarbonate concentration. According to

prior art, this problem is solved either by use of a gas barrier for carbon dioxide or by allowing the bicarbonate to slowly equilibrate with the atmosphere.

If a gas barrier is used, a complicated and  
5 expensive polymer is required as gas barrier otherwise it will result, after mixing with the rest of the content in the container, in an non-definable pH (depending of age of the solution). The polymers used for these types of gas barriers are often very brittle, and care has to be  
10 taken when handling and storing the bags with gas barriers not to create cracks that will give rise to leakage.

The idea of letting the bicarbonate slowly equilibrate with the atmosphere is for example disclosed  
15 in US 6,309,673. However, this way of solving the problem creates an uncertainty concerning the pH value and the bicarbonate concentration in the final, ready-to-use solution.

In US 5,296,242 a solution is disclosed in which a  
20 premix of bicarbonate and carbonate is used in the buffer system. This document discloses a specific mix of bicarbonate and carbonate, which provides for a partial pressure of carbon dioxide that equals the physiological value of the partial pressure within the human body. The  
25 buffer solution is further combined with an acid solution, the acid being a metabolizable, organic acid. This reference stresses that an organic acid should be used, this for the therapy of acidosis.

### 30 Summary of the invention

An object of the invention is to provide a medical solution that, on one hand, ensures good stability, and on the other hand, ensures good biocompatibility.

The present invention provides a medical solution comprising at least two single solutions which, after terminal sterilization and up on use, can be mixed and used as a medical solution. The first single solution  
5 comprises bicarbonate and carbonate in such proportions that the partial pressure of carbon dioxide,  $\text{CO}_2$ , in the first single solution is of the same order of magnitude as the partial pressure of carbon dioxide,  $\text{CO}_2$ , of the atmosphere. The second single solution comprises an acid  
10 and has a pH of 1,0-1,5. When said first and second single solutions, after terminal sterilization and up on use, are mixed a final solution, ready for use, is formed and it has a pH within the range of 7.0-7.6.

By ensuring that the partial pressure of carbon  
15 dioxide,  $\text{CO}_2$ , in the first single solution is of the same order of magnitude as the partial pressure of carbon dioxide,  $\text{CO}_2$ , of the atmosphere, there is no driving force which urges carbon dioxide out from the container in which the solutions are contained. Accordingly, the  
20 carbonate and bicarbonate concentrations will be stable and the problem with the solutions of prior art is overcome.

In a preferred embodiment of the present invention said first single solution has a pH of 10.1-10.5,  
25 preferably 10.3.

Said second single solution preferably has a pH of 1.3. Preferably, said second single solution is acidified by hydrochloric acid,  $\text{HCl}$ .

According to another preferred embodiment of the  
30 present invention the medical solution further comprises one or more osmotic agents. Preferred said one or more osmotic agents are chosen among glucose, glucose polymers, glycerol, xylitol, fructose, amino acids,

peptides, proteins, amino sugars, N-acetyl glucose amine (NAG), or combinations thereof.

In one preferred embodiment of the present invention said one or more osmotic agents are, before being mixed  
5 into the final solution, arranged in said second single solution. In another preferred embodiment, said one or more osmotic agents are arranged in a third single solution before being mixed into the final solution. In even another preferred embodiment, said one or more  
10 osmotic agents are also arranged in a fourth single solution before being mixed into the final solution.

In a preferred embodiment, said one or more osmotic agents in said third and fourth single solutions is glucose and/or glucose polymers, which could give rise to  
15 glucose degradation products (GDPs) during terminal sterilization and/or storage. If this is the case, said third and fourth single solutions comprise an acid and have a pH of at least 1.8, preferably at least 2.0, and a pH of at most 2.6, preferably at most 2.5, and most  
20 preferably at most 2.3.

According to another preferred embodiment of the present invention the final solution further comprises one or more electrolytes. The one or more electrolytes comprise according to one preferred embodiment of the  
25 invention one or more of the ions of sodium, calcium, potassium, magnesium and/or chloride. These one or more electrolytes could, according to different preferred embodiments of the present invention, before being mixed into the final solution, be included in said first single  
30 solution, in said second single solution, and/or in said optional third and/or fourth single solution. However, magnesium and/or calcium should not be included in said first single solution, as magnesium and calcium will

precipitate as magnesium carbonate and calcium carbonate, respectively, when included in the first single solution together with carbonate and bicarbonate.

In another preferred embodiment of the invention,  
5 the different single solutions are provided in different compartments in a multi-compartment bag before being mixed to the final solution.

The present invention further provides for a method for producing said medical solution. According to the  
10 invention the method comprises providing said single solutions in separate compartments, and thereafter terminally sterilizing said single solutions.

According to a preferred embodiment of the method according to the present invention said terminal  
15 sterilization is heat sterilization and/or radiation sterilization. In an even more preferred embodiment of the method according to the present invention, said terminal sterilization is heat sterilization at a temperature of at least 100°C, preferably at least 121 °C.

20 In a preferred embodiment of the invention said first and second single solutions, after sterilization and up on use, are mixed to form a final solution. In another preferred embodiment said first, second and third single solutions, after sterilization and up on use, are  
25 mixed to form a final solution. In another preferred embodiment of the invention, said first, second and fourth single solutions, after sterilization and up on use, are mixed to form a final solution, and in even another preferred embodiment of the invention, said  
30 first, second, third and fourth single solutions, after sterilization and up on use, are mixed to form a final solution.

The present invention further provides a multi-compartment bag comprising the medical solution according to above and the use of the medical solution according to above.

5 Additional objects, features, advantages and preferred embodiments of the present invention will become apparent from the following detailed description when taken in conjunction with the enclosed patent claims.

10

#### Definitions

The term "medical solution" is intended to mean dialysis solutions for hemodialysis, hemodiafiltration, hemofiltration, and peritoneal dialysis, solutions for  
15 dialysis within renal intensive care, solutions for substitution or infusion normally containing buffering substances, and solutions for nutrition purposes.

The term "single solution" is intended to mean one solution kept isolated from other solutions up until use.

20 The term "bicarbonate and carbonate" is intended to mean alkali bicarbonate and alkali carbonate, especially sodium bicarbonate and sodium carbonate.

The term "a final solution" is intended to mean the solution which includes the required different single  
25 solutions and which is ready for use.

The term "multi-compartment bag" is intended to mean bag divided into more than one compartment and that the content in the different compartments could be brought together and mixed before use.

30 The term "terminal sterilization" is intended to mean that the product is sterilized in its final package. The terminal sterilization may include heat sterilization and/or radiation sterilization, but is preferably heat

sterilization effected in an autoclave at a temperature of at least 100°C, preferably at least 121°C.

The term "up on use" is intended to mean as close as possible before the medical solution is used for its  
5 specific purpose.

#### Detailed description of the invention

The medical solution according to the invention comprises a first single solution and a second single  
10 solution, wherein said first and second single solutions, after terminal sterilization and up on use, are to be mixed to form a final solution.

Said first single solution comprises bicarbonate and carbonate in such proportions that the partial pressure  
15 of carbon dioxide, CO<sub>2</sub>, in the first single solution is of the same order of magnitude as the partial pressure of carbon dioxide, CO<sub>2</sub>, of the atmosphere. Preferably, bicarbonate and carbonate are mixed as sodium bicarbonate and sodium carbonate. Preferably, said first single  
20 solution has a pH within the range of 10.1-10.5, most preferably said first single solution has a pH of 10.3.

After having mixed at least said first and second single solutions into a final solution, said final solution has a pH within the range of 7.0-7.6. Further,  
25 said final solution preferably has a bicarbonate concentration of at least 25 mM, preferably at least 30 mM, and at most 45 mM, preferably at most 40 mM.

Said second single solution has preferably a pH within the range of 1.0-1.5, most preferably a pH of 1.3.  
30 In a preferred embodiment of the invention said second single solution comprises HCl.

The medical solution according to the invention preferably comprises one or more osmotic agents that are

preferably chosen among glucose, glucose polymers, glycerol, xylitol, fructose, amino acids, peptides, proteins, amino sugars, N-acetyl glucose amine (NAG), or combinations thereof. The one or more osmotic agents are  
5 in one preferred embodiment, before being mixed into the final solution, arranged in said second single solution. However, in another preferred embodiment the one or more osmotic agents are arranged in a third single solution. In even another preferred embodiment of the invention,  
10 said one or more osmotic agents are, before being mixed into said final solution besides being arranged in a third single solution, also arranged in a fourth single solution.

In case of using one or more osmotic agents, which  
15 could give rise to glucose degradation products, said third and fourth single solutions further comprise an acid and preferably have a pH of at least 1.8, preferably at least 2.0, and a pH of at most 2.6, preferably at most 2.5 and most preferably at most 2.3. Within these pH  
20 ranges the amount of the glucose degradation products (GDPs) being most toxic is as low as possible, and especially 3,4-dideoxyglucosone-3-ene (3,4-DGE), which is the most toxic one of all the GDPs. GDPs are known to give rise to several problems during for example  
25 peritoneal dialysis and of course it is always an aim to reduce the amount of toxic substances. However, besides optimizing the pH of said third and forth single solutions, it is also important to keep the concentration of one or more osmotic agents, which could give rise to  
30 GDPs, of at least 10 % by weight, preferably at least 20 % by weight and most preferably at least 40 % by weight, based on the total weight of said third and fourth single solutions, respectively.



In a preferred embodiment of the invention said third and fourth single solutions could comprise different total amounts of one or more osmotic agents. The different total amounts could be achieved by

5 providing the same concentrations within said third and fourth single solutions, but providing different volumes thereof. The different total amounts could also be achieved by providing the same volume of said third and fourth single solutions, but providing different

10 concentrations in said third single solution in comparison with said fourth single solution. By having such a preferred medical solution comprising said first, second, third and fourth single solutions, the user thereof could choose what concentration of osmotic agent

15 the user would like to have for a specific treatment. By combining said first, second and third single solutions to a final solution, the user gets a first specific concentration of osmotic agent, by combining said first, second and fourth single solutions to a final solution,

20 the user gets a second specific concentration of osmotic agent, and by combining said first, second, third and fourth single solutions to a final solution, the user gets a third specific concentration of osmotic agent. Accordingly, said third and fourth single solutions

25 could, up on use, be mixed individually, with said first and second solutions, i.e. either first, second and third, or first, second and fourth, or jointly, i.e. mixing first, second, third and fourth single solutions together. Note that said final solution always have a pH

30 within the range of 7.0-7.6, no matter which of the above combinations of single solutions is used. The buffer solution in said first single solution have the capability to buffer said third and/or fourth single

solution(s) in combination with said second single solution to a final solution with a pH of 7.0-7.6.

In a preferred embodiment of the invention the medical solution further contains one or more  
5 electrolytes. Preferably, the electrolytes is one or more of the ions of sodium, calcium, potassium, magnesium and chloride.

The arrangement of electrolytes in the different compartments is dependent on the different electrolytes  
10 co-behavior with the other substances present in the single solutions, i.e. whether some sort of reaction could occur between one or more of the electrolyte(s) and the other substances present in a specific single solution. Usually, the electrolytes are contained in said  
15 second single solution. For example, calcium and magnesium ions are preferably provided in any of the other single solutions, but said first single solution. The reason for this is that calcium and magnesium together with bicarbonate and/or carbonate could cause  
20 precipitation of calcium carbonate and magnesium carbonate, respectively. However, calcium ions could be kept with bicarbonate under certain circumstances, such as specific pH ranges and so on, this is for example disclosed in EP 0 437 274, which hereby is enclosed by  
25 reference.

In the method for producing a medical solution according to above, said single solutions are provided in separate compartments. Thereafter said single solutions are terminally sterilized. Preferably, the terminal  
30 sterilization is heat sterilization and/or radiation sterilization, (see also European Pharmacopoeia 1977 for a review of different sterilization techniques). In a preferred embodiment of the method according to the

invention, the terminal sterilization is heat sterilization at a temperature of at least 100°C, preferably at least 121°C.

The sterilization time may vary depending on the  
5 sterilization temperature, the type of container and the contents therein to be sterilized.

The radiation sterilization may be either ionising or non-ionising sterilization. Examples of ionising sterilization are gamma and beta radiation. Examples of  
10 non-ionising radiation sterilization is UV radiation.

The medical solution according to the present invention has the advantage of ensuring good stability and good biocompatibility.

Said single solutions could be provided in different  
15 compartments in a multi-compartment bag, and the mixing could be provided by having the different compartments sealingly coupled by frangible pins, which different pins could be broken in order to mix the content in optional compartments within the multicompartment bag. The mixing  
20 could further be provided by having a peel seal in-between the different compartments, which peel seals could be peeled in order to mix the content in the different compartments.

Below you will find different examples of solutions  
25 according to the present invention.

#### Exampels

By way of example, and not limitation, the following examples identify a variety of solutions made pursuant to  
30 an embodiment of the present invention.

Example 1 - 4

Two compartment bags

Compartment 1:

	Example 1	Example 2	Example 3	Example 4
NaHCO <sub>3</sub>	95,5 mM	95,5 mM	112 mM	112 mM
Na <sub>2</sub> CO <sub>3</sub>	304,5 mM	304,5 mM	258 mM	258 mM
pH	10,3	10,3	10,3	10,3
Volume	0,5 l	0,5 l	0,5 l	0,5 l

5

Compartment 2:

	Example 1	Example 2	Example 3	Example 4
HCl	38 mM	38 mM	38 mM	38 mM
NaCl	77,3 mM	77,3 mM	77,3 mM	77,3 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,95 mM	1,95 mM	1,95 mM	1,95 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,56 mM	0,56 mM	0,56 mM	0,56 mM
Glucose	-	1,22 g/l	-	1,22 g/l
Lactate	-	-	3,30 mM	3,30 mM
pH	1,3	1,3	1,3	1,3
Volume	4,5 l	4,5 l	4,5 l	4,5 l

Solution, mixed and ready for use:

	Example 1	Example 2	Example 3	Example 4
Volume	5 l	5 l	5 l	5 l
pH	7,25	7,25	7,25	7,25
Cl <sup>-</sup>	108,3 mM	108,3 mM	108,3 mM	108,3 mM
Na <sup>+</sup>	140,02 mM	140,02 mM	135,37 mM	135,37 mM
Ca <sup>+</sup>	1,76 mM	1,76 mM	1,76 mM	1,76 mM
Mg <sup>+</sup>	0,5 mM	0,5 mM	0,5 mM	0,5 mM
HCO <sub>3</sub> <sup>-</sup>	40 mM	40 mM	37 mM	37 mM
Glucose	-	1,1 g/l	-	1,1 g/l
Lactate	-	-	3 mM	3 mM

Example 5-8

Two compartment bags

Compartment 1:

	Example 5	Example 6	Example 7	Example 8
NaHCO <sub>3</sub>	139 mM	139 mM	133 mM	133 mM
Na <sub>2</sub> CO <sub>3</sub>	661 mM	661 mM	607 mM	607 mM
pH	10,3	10,3	10,3	10,3
Volume	0,25 l	0,25 l	0,25 l	0,25 l

5

Compartment 2:

	Example 5	Example 6	Example 7	Example 8
HCl	38,4 mM	38,4 mM	38,4 mM	38,4 mM
NaCl	70,5 mM	70,5 mM	70,5 mM	70,5 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,84 mM	1,84 mM	1,84 mM	1,84 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,53 mM	0,53 mM	0,53 mM	0,53 mM
Glucose	-	1,16 g/l	-	1,16 g/l
Lactate	-	-	3,16 mM	3,16 mM
pH	1,3	1,3	1,3	1,3
Volume	4,75 l	4,75 l	4,75 l	4,75 l

Solution, mixed and ready for use:

	Example 5	Example 6	Example 7	Example 8
Volume	5 l	5 l	5 l	5 l
pH	7,25	7,25	7,25	7,25
Cl <sup>-</sup>	107,6 mM	107,6 mM	107,6 mM	107,6 mM
Na <sup>+</sup>	140,03 mM	140,03 mM	137,18 mM	137,18 mM
Ca <sup>+</sup>	1,75 mM	1,75 mM	1,75 mM	1,75 mM
Mg <sup>+</sup>	0,5 mM	0,5 mM	0,5 mM	0,5 mM
HCO <sub>3</sub> <sup>-</sup>	40 mM	40 mM	37 mM	37 mM
Glucose	-	1,1 g/l	-	1,1 g/l
Lactate	-	-	3 mM	3 mM

Examples 9-12

Three compartment bags:

Compartment 1:

	Example 9	Example 10	Example 11	Example 12
NaHCO <sub>3</sub>	95,5 mM	112 mM	139 mM	133 mM
Na <sub>2</sub> CO <sub>3</sub>	304,5 mM	258 mM	661 mM	607 mM
pH	10,3	10,3	10,3	10,3
Volume	0,2 l	0,2 l	0,1 l	0,1 l

5

Compartment 2:

	Example 9	Example 10	Example 11	Example 12
HCl	41,02 mM	41,02 mM	39,23 mM	39,23mM
NaCl	83,45 mM	83,45 mM	72,02 mM	72,02 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	2,11 mM	2,11 mM	1,88 mM	1,88 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,60 mM	0,60 mM	0,54 mM	0,54 mM
Glucose	17,05	17,05 g/l	16,13 g/l	16,13 g/l
Lactate	-	3,41 mM	-	3,23 mM
pH	1,3	1,3	1,3	1,3
Volume	1,76 l	1,76 l	1,86 l	1,86 l

Third compartment:

	Example 9	Example 10	Example 11	Example 12
Glucose	500 g/l	500 g/l mM	500 g/lmM	500 g/l
pH	2-2,5	2-2,5	2-2,5	2-2,5
Volume	0,04 l	0,04 l	0,04 l	0,04 l

15

Solution, mixed and ready for use:

	Example 9	Example 10	Example 11	Example 12
Volume	2 l	2 l	2 l	2 l
pH	7,25	7,25	7,25	7,25
Cl <sup>-</sup>	108,3 mM	108,3 mM	107,96 mM	107,96 mM
Na <sup>+</sup>	140,02 mM	135,37 mM	140,03 mM	140,03 mM
Ca <sup>+</sup>	1,76 mM	1,76 mM	1,75 mM	1,75 mM
Mg <sup>+</sup>	0,5 mM	0,5 mM	0,5 mM	0,5 mM
HCO <sub>3</sub> <sup>-</sup>	40 mM	37 mM	40 mM	40 mM
Glucose	15/25 g/l*	15/25 g/l*	15/25 g/l*	15/25 g/l*
Lactate	-	3 mM	-	3 mM

\* 15 g/l is compartment 1 and 2 are mixed and 25 g/l if all three compartments are mixed.

5

Examples 13-16

Four-compartment bags:

10 Compartment 1:

	Example 13	Example 14	Example 15	Example 16
NaHCO <sub>3</sub>	95,5 mM	112 mM	139 mM	133 mM
Na <sub>2</sub> CO <sub>3</sub>	304,5 mM	258 mM	661 mM	607 mM
pH	10,3	10,3	10,3	10,3
Volume	0,196 l	0,196 l	0,098 l	0,098 l

16

Compartment 2:

	Example 13	Example 14	Example 15	Example 16
HCl	38,78 mM	38,78 mM	39,18 mM	39,18 mM
NaCl	78,88 mM	78,88 mM	71,94 mM	71,94 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,99 mM	1,99 mM	1,88 mM	1,88 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,57 mM	0,57 mM	0,54 mM	0,54 mM
Lactate	-	3,41 mM	-	3,22 mM
pH	1,3	1,3	1,3	1,3
Volume	1,764 l	1,764 l	1,862 l	1,862 l

Third compartment:

	Example 13	Example 14	Example 15	Example 16
Glucose	500 g/l	500 g/l	500 g/l	500 g/l
pH	2-2,5	2-2,5	2-2,5	2-2,5
Volume	0,062 l	0,062 l	0,062 l	0,062 l

5 Fourth compartment:

	Example 13	Example 14	Example 15	Example 16
Glucose	500 g/l	500 g/l	500 g/l	500 g/l
pH	2-2,5	2-2,5	2-2,5	2-2,5
Volume	0,103 l	0,103 l	0,103 l	0,103 l



17

Solution, mixed and ready for use:

Example 13	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,25	7,25	7,25
Cl <sup>-</sup>	107,1 mM	105,0 mM	101,9 mM
Na <sup>+</sup>	137,1 mM	134,4 mM	130,5 mM
Ca <sup>+</sup>	1,74 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l

Example 14	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,25	7,25	7,25
Cl <sup>-</sup>	107,1 mM	105,0 mM	101,9 mM
Na <sup>+</sup>	135,6 mM	132,9 mM	129,1 mM
Ca <sup>+</sup>	1,74 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	35,9 mM	35,2 mM	34,1 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l
Lactate	2,98 mM	2,92 mM	2,83 mM

18

Example 15	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,25	7,25	7,25
Cl <sup>-</sup>	106,8 mM	104,7 mM	101,6 mM
Na <sup>+</sup>	137,1 mM	134,3 mM	130,5 mM
Ca <sup>+</sup>	1,73 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l

Example 16	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,25	7,25	7,25
Cl <sup>-</sup>	106,8 mM	104,7 mM	101,6 mM
Na <sup>+</sup>	137,5 mM	134,7 mM	130,8 mM
Ca <sup>+</sup>	1,73 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,5 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l
Lactate	2,97 mM	2,91 mM	2,82 mM

Example 17-20

## 5 Two compartment bags

## Compartment 1:

	Example 17	Example 18	Example 19	Example 20
NaHCO <sub>3</sub>	95,5 mM	95,5 mM	112 mM	112 mM
Na <sub>2</sub> CO <sub>3</sub>	304,5 mM	304,5 mM	258 mM	258 mM
pH	10,3	10,3	10,3	10,3
Volume	0,5 l	0,5 l	0,5 l	0,5 l

19

Compartment 2:

	Example 17	Example 18	Example 19	Example 20
HCl	35,5 mM	35,5 mM	35,5 mM	35,5 mM
NaCl	77,3 mM	77,3 mM	77,3 mM	77,3 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,95 mM	1,95 mM	1,95 mM	1,95 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,56 mM	0,56 mM	0,56 mM	0,56 mM
Glucose	-	1,22 g/l	-	1,22 g/l
Lactate	-	-	3,30 mM	3,30 mM
pH	1,3	1,3	1,3	1,3
Volume	4,5 l	4,5 l	4,5 l	4,5 l

Solution, mixed and ready for use:

	Example 17	Example 18	Example 19	Example 20
Volume	5 l	5 l	5 l	5 l
pH	7,5	7,5	7,5	7,5
Cl <sup>-</sup>	106,03 mM	106,03 mM	106,03 mM	106,03 mM
Na <sup>+</sup>	140,02 mM	140,02 mM	135,37 mM	135,37 mM
Ca <sup>+</sup>	1,76 mM	1,76 mM	1,76 mM	1,76 mM
Mg <sup>+</sup>	0,5 mM	0,5 mM	0,5 mM	0,5 mM
HCO <sub>3</sub> <sup>-</sup>	40 mM	40 mM	37 mM	37 mM
Glucose	-	1,1 g/l	-	1,1 g/l
Lactate	-	-	3 mM	3 mM

5 Example 21-24

Two compartment bags

Compartment 1:

	Example 21	Example 22	Example 23	Example 24
NaHCO <sub>3</sub>	139 mM	139 mM	133 mM	133 mM
Na <sub>2</sub> CO <sub>3</sub>	661 mM	661 mM	607 mM	607 mM
pH	10,3	10,3	10,3	10,3
Volume	0,25 l	0,25 l	0,25 l	0,25 l

## Compartment 2:

	Example 21	Example 22	Example 23	Example 24
HCl	36,11 mM	36,11 mM	36,11 mM	36,11 mM
NaCl	70,5 mM	70,5 mM	70,5 mM	70,5 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,84 mM	1,84 mM	1,84 mM	1,84 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,53 mM	0,53 mM	0,53 mM	0,53 mM
Glucose	-	1,16 g/l	-	1,16 g/l
Lactate	-	-	3,16 mM	3,16 mM
pH	1,3	1,3	1,3	1,3
Volume	4,75 l	4,75 l	4,75 l	4,75 l

## Solution, mixed and ready for use:

	Example 21	Example 22	Example 23	Example 24
Volume	5 l	5 l	5 l	5 l
pH	7,5	7,5	7,5	7,5
Cl <sup>-</sup>	105,78 mM	105,78 mM	105,78 mM	105,78 mM
Na <sup>+</sup>	140,03 mM	140,03 mM	137,18 mM	137,18 mM
Ca <sup>+</sup>	1,75 mM	1,75 mM	1,75 mM	1,75 mM
Mg <sup>+</sup>	0,5 mM	0,5 mM	0,5 mM	0,5 mM
HCO <sub>3</sub> <sup>-</sup>	40 mM	40 mM	37 mM	37 mM
Glucose	-	1,1 g/l	-	1,1 g/l
Lactate	-	-	3 mM	3 mM

Example 25-28

Two compartment bags

Compartment 1:

	Example 25	Example 26	Example 27	Example 28
NaHCO <sub>3</sub>	95,5 mM	95,5 mM	112 mM	112 mM
Na <sub>2</sub> CO <sub>3</sub>	304,5 mM	304,5 mM	258 mM	258 mM
pH	10,3	10,3	10,3	10,3
Volume	0,5 l	0,5 l	0,5 l	0,5 l

5

Compartment 2:

	Example 25	Example 26	Example 27	Example 28
HCl	38,74 mM	38,74 mM	38,74 mM	38,74 mM
NaCl	77,3 mM	77,3 mM	77,3 mM	77,3 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,95 mM	1,95 mM	1,95 mM	1,95 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,56 mM	0,56 mM	0,56 mM	0,56 mM
Glucose	-	1,22 g/l	-	1,22 g/l
Lactate	-	-	3,30 mM	3,30 mM
pH	1,3	1,3	1,3	1,3
Volume	4,5 l	4,5 l	4,5 l	4,5 l

Solution, mixed and ready for use:

	Example 25	Example 26	Example 27	Example 28
Volume	5 l	5 l	5 l	5 l
pH	7,0	7,0	7,0	7,0
Cl <sup>-</sup>	108,95 mM	108,95 mM	108,95 mM	108,95 mM
Na <sup>+</sup>	140,02 mM	140,02 mM	135,37 mM	135,37 mM
Ca <sup>+</sup>	1,76 mM	1,76 mM	1,76 mM	1,76 mM
Mg <sup>+</sup>	0,5 mM	0,5 mM	0,5 mM	0,5 mM
HCO <sub>3</sub> <sup>-</sup>	40 mM	40 mM	37 mM	37 mM
Glucose	-	1,1 g/l	-	1,1 g/l
Lactate	-	-	3 mM	3 mM

Example 29-32

Two compartment bags

## 5 Compartment 1:

	Example 29	Example 30	Example 31	Example 32
NaHCO <sub>3</sub>	139 mM	139 mM	133 mM	133 mM
Na <sub>2</sub> CO <sub>3</sub>	661 mM	661 mM	607 mM	607 mM
pH	10,3	10,3	10,3	10,3
Volume	0,25 l	0,25 l	0,25 l	0,25 l

## Compartment 2:

	Example 29	Example 30	Example 31	Example 32
HCl	39,19 mM	39,19 mM	39,19 mM	39,19 mM
NaCl	70,5 mM	70,5 mM	70,5 mM	70,5 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,84 mM	1,84 mM	1,84 mM	1,84 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,53 mM	0,53 mM	0,53 mM	0,53 mM
Glucose	-	1,16 g/l	-	1,16 g/l
Lactate	-	-	3,16 mM	3,16 mM
pH	1,3	1,3	1,3	1,3
Volume	4,75 l	4,75 l	4,75 l	4,75 l

23

Solution, mixed and ready for use:

	Example 29	Example 30	Example 31	Example 32
Volume	5 l	5 l	5 l	5 l
pH	7,0	7,0	7,0	7,0
Cl <sup>-</sup>	108,71 mM	108,71 mM	108,71 mM	108,71 mM
Na <sup>+</sup>	140,03 mM	140,03 mM	137,18 mM	137,18 mM
Ca <sup>+</sup>	1,75 mM	1,75 mM	1,75 mM	1,75 mM
Mg <sup>+</sup>	0,5 mM	0,5 mM	0,5 mM	0,5 mM
HCO <sub>3</sub> <sup>-</sup>	40 mM	40 mM	37 mM	37 mM
Glucose	-	1,1 g/l	-	1,1 g/l
Lactate	-	-	3 mM	3 mM

Examples 33-36

## 5 Four-compartment bags:

## Compartment 1:

	Example 33	Example 34	Example 35	Example 36
NaHCO <sub>3</sub>	95,5 mM	112 mM	139 mM	133 mM
Na <sub>2</sub> CO <sub>3</sub>	304,5 mM	258 mM	661 mM	607 mM
pH	10,3	10,3	10,3	10,3
Volume	0,196 l	0,196 l	0,098 l	0,098 l

## Compartment 2:

	Example 33	Example 34	Example 35	Example 36
HCl	38,74 mM	38,74 mM	39,19 mM	39,19 mM
NaCl	78,88 mM	78,88 mM	71,94 mM	71,94 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,99 mM	1,99 mM	1,88 mM	1,88 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,57 mM	0,57 mM	0,54 mM	0,54 mM
Lactate	-	3,41 mM	-	3,22 mM
pH	1,3	1,3	1,3	1,3
Volume	1,764 l	1,764 l	1,862 l	1,862 l

Third compartment:

	Example 33	Example 34	Example 35	Example 36
Glucose	500 g/l	500 g/l	500 g/l	500 g/l
pH	2-2,5	2-2,5	2-2,5	2-2,5
Volume	0,062 l	0,062 l	0,062 l	0,062 l

Fourth compartment:

	Example 33	Example 34	Example 35	Example 36
Glucose	500 g/l	500 g/l	500 g/l	500 g/l
pH	2-2,5	2-2,5	2-2,5	2-2,5
Volume	0,103 l	0,103 l	0,103 l	0,103 l

5 Solution, mixed and ready for use:

Example 33	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,0	7,0	7,0
Cl <sup>-</sup>	107,8 mM	105,6 mM	102,5 mM
Na <sup>+</sup>	137,1 mM	134,4 mM	130,5 mM
Ca <sup>+</sup>	1,74 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l



25

Example 34	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,0	7,0	7,0
Cl <sup>-</sup>	107,8 mM	105,6 mM	102,5 mM
Na <sup>+</sup>	135,6 mM	132,9 mM	129,1 mM
Ca <sup>+</sup>	1,74 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	35,9 mM	35,2 mM	34,1 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l
Lactate	2,98 mM	2,92 mM	2,83 mM

Example 35	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,0	7,0	7,0
Cl <sup>-</sup>	113,2 mM	110,9mM	107,7 mM
Na <sup>+</sup>	137,1 mM	134,3 mM	130,5 mM
Ca <sup>+</sup>	1,73 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l

26

Example 36	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,0	7,0	7,0
Cl <sup>-</sup>	113,2 mM	110,9 mM	107,7 mM
Na <sup>+</sup>	137,5 mM	134,7 mM	130,8 mM
Ca <sup>+</sup>	1,73 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,5 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l
Lactate	2,97 mM	2,91 mM	2,82 mM

Examples 37-40

Four-compartment bags:

5

Compartment 1:

	Example 37	Example 38	Example 39	Example 40
NaHCO <sub>3</sub>	95,5 mM	112 mM	139 mM	133 mM
Na <sub>2</sub> CO <sub>3</sub>	304,5 mM	258 mM	661 mM	607 mM
pH	10,3	10,3	10,3	10,3
Volume	0,196 l	0,196 l	0,098 l	0,098 l

Compartment 2:

	Example 37	Example 38	Example 39	Example 40
HCl	35,49 mM	35,49 mM	36,11 mM	36,11 mM
NaCl	78,88 mM	78,88 mM	71,94 mM	71,94 mM
CaCl <sub>2</sub> *2H <sub>2</sub> O	1,99 mM	1,99 mM	1,88 mM	1,88 mM
MgCl <sub>2</sub> *6 H <sub>2</sub> O	0,57 mM	0,57 mM	0,54 mM	0,54 mM
Lactate	-	3,41 mM	-	3,22 mM
pH	1,3	1,3	1,3	1,3
Volume	1,764 l	1,764 l	1,862 l	1,862 l

Third compartment:

	Example 37	Example 38	Example 39	Example 40
Glucose	500 g/l	500 g/l	500 g/l	500 g/l
pH	2-2,5	2-2,5	2-2,5	2-2,5
Volume	0,062 l	0,062 l	0,062 l	0,062 l

Fourth compartment:

	Example 37	Example 38	Example 39	Example 40
Glucose	500 g/l	500 g/l	500 g/l	500 g/l
pH	2-2,5	2-2,5	2-2,5	2-2,5
Volume	0,103 l	0,103 l	0,103 l	0,103 l

5 Solution, mixed and ready for use:

Example 37	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,5	7,5	7,5
Cl <sup>-</sup>	104,9 mM	102,8 mM	99,8 mM
Na <sup>+</sup>	137,1 mM	134,4 mM	130,5 mM
Ca <sup>+</sup>	1,74 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l

28

Example	Compartment	Compartment	Compartment
38	1+2+3	1+2+4	1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,5	7,5	7,5
Cl <sup>-</sup>	104,9 mM	102,8 mM	99,8 mM
Na <sup>+</sup>	135,6 mM	132,9 mM	129,1 mM
Ca <sup>+</sup>	1,74 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	35,9 mM	35,2 mM	34,1 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l
Lactate	2,98 mM	2,92 mM	2,83 mM

Example	Compartment	Compartment	Compartment
39	1+2+3	1+2+4	1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,5	7,5	7,5
Cl <sup>-</sup>	110,1 mM	107,9 mM	104,8 mM
Na <sup>+</sup>	137,1 mM	134,3 mM	130,5 mM
Ca <sup>+</sup>	1,73 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,50 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l

Example 40	Compartment 1+2+3	Compartment 1+2+4	Compartment 1+2+3+4
Volume	2,022 l	2,063 l	2,125 l
pH	7,5	7,5	7,5
Cl <sup>-</sup>	110,1 mM	107,9 mM	104,8 mM
Na <sup>+</sup>	137,5 mM	134,7 mM	130,8 mM
Ca <sup>+</sup>	1,73 mM	1,70 mM	1,65 mM
Mg <sup>+</sup>	0,5 mM	0,49 mM	0,47 mM
HCO <sub>3</sub> <sup>-</sup>	38,8 mM	38,0 mM	36,9 mM
Glucose	15,3 g/l	25,0 g/l	38,8 g/l
Lactate	2,97 mM	2,91 mM	2,82 mM

In summary, based on the above results, the inventors concluded that a stable and biocompatible bicarbonate-based solution can be prepared, provided that it comprises bicarbonate and carbonate in such proportions that the partial pressure of carbon dioxide, CO<sub>2</sub>, is of the same order of magnitude as the partial pressure of CO<sub>2</sub> of the atmosphere.

It should be understood that various changes and modifications to the presently preferred embodiments described herein will be apparent to those skilled in the art. Such changes and modifications can be made without departing from the spirit and scope of the present invention and without diminishing its attendant advantages. It is therefore intended that such changes and modifications be covered by the appended claims.